



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 43/90, 43/653 // (A01N 43/90, 43:653, 43:56, 43:42) (A01N 43/653, 43:653, 43:56, 43:42)		A1	(11) International Publication Number: WO 99/57982 (43) International Publication Date: 18 November 1999 (18.11.99)
(21) International Application Number: PCT/US99/10359 (22) International Filing Date: 12 May 1999 (12.05.99) (30) Priority Data: 60/085,397 14 May 1998 (14.05.98) US 60/122,791 4 March 1999 (04.03.99) US (71) Applicant (for all designated States except US): E.I. DU PONT DE NEMOURS AND COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): LEEP, Daniel, Carl [US/US]; 12 Farmingdale Lane, Newark, DE 19711 (US). (74) Agent: BIRCH, Linda, D.; E.I. du Pont de Nemours and Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).		(81) Designated States: AE, AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(54) Title: CROP-SAFENED HERBICIDAL MIXTURES			
<div style="display: flex; justify-content: space-around; align-items: flex-start;"> <div style="text-align: center;"> <p>(Ia)</p> </div> <div style="text-align: center;"> <p>(Ib)</p> </div> </div> <div style="display: flex; justify-content: space-around; align-items: flex-start; margin-top: 20px;"> <div style="text-align: center;"> <p>(IIa)</p> </div> <div style="text-align: center;"> <p>(IIb)</p> </div> </div> <div style="text-align: center; margin-top: 20px;"> <p>(IIc)</p> </div>			
(57) Abstract			
<p>This invention relates to herbicidal mixtures of at least one compound selected from Formulae (Ia) and (Ib) with a safener selected from Formulae (IIa), (IIb) and (IIc) wherein R¹, R², R³ and R⁴ are as defined in the text, herbicidal compositions of said mixtures, and a method for the use of said mixtures to control undesired vegetation.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

TITLE

CROP-SAFENED HERBICIDAL MIXTURES

FIELD OF THE INVENTION

The present invention relates to mixtures of herbicides with safeners that reduce crop
 5 injury while retaining weed control.


BACKGROUND OF THE INVENTION

The control of undesired vegetation is extremely important in achieving high crop
 efficiency. This can be achieved by the selective control of the growth of weeds in such
 useful crops as alfalfa, corn (maize), peanut (groundnut), potato, rape, rice, sorghum,
 10 soybean, sugar beet, tomato, vegetables, cool-season cereals including wheat, barley and rye,
 perennial plantation crops including banana, citrus, cocoa, coffee, grapes, hops, plantain,
 pineapple, oil palm, rubber, sugarcane, tea, fruit trees, nut trees and forests, and turf, among
 others. Unchecked weed growth in such useful crops can cause significant reduction in
 productivity and thereby result in increased costs to the consumers. The need for finding
 15 products that achieve such results continues to be commercially important.


Herbicidally active compounds are typically applied in combination with nonherbicidal
 formulating ingredients that facilitate application, increase wash-off resistance, improve
 penetration, etc. Certain rare combinations of herbicides with nonherbicidal compounds
 surprisingly reduce the herbicidal effect on crops while retaining useful weed control. The
 20 nonherbicidal compounds having this effect are termed "safeners" or "antidotes". By
 reducing crop injury while retaining control of weeds, effective weed-control selectivity is
 increased. Such valuable combinations have now been discovered.

SUMMARY OF THE INVENTION


This invention relates to mixtures of chlorofluorobenzene herbicides with several
 25 compounds discovered to be useful safeners for these herbicides. The herbicidal mixtures
 comprise herbicidally effective amounts of at least one herbicidal compound selected from
 the group consisting of:

**1a**

(6*S*-*cis*)-1-chloro-*N*-[2-chloro-4-fluoro-5-(6-fluorotetrahydro-1,3-dioxo-1*H*-pyrrolo-
 30 [1,2-*c*]imidazol-2(3*H*)-yl)phenyl]methanesulfonamide (**1a**), or an agriculturally useful salt
 thereof; and


**Ib**

ethyl $\alpha,2$ -dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1*H*-1,2,4-triazol-1-yl]-4-fluorobenzenepropanoate (carfentrazone-ethyl, **Ib**), or an agriculturally useful salt thereof; in combination with an antidotally effective amount of a safener compound selected from the group consisting of:

**IIa**

or an agriculturally useful salt thereof
wherein

R^1 is C_1 - C_{12} alkyl, C_3 - C_7 cycloalkyl, C_2 - C_8 alkenyl or C_2 - C_8 alkynyl, each optionally substituted with at least one radical selected from the group consisting of halogen, hydroxy, C_1 - C_4 alkoxy and C_1 - C_4 alkylthio;


**IIb**

or an agriculturally useful salt thereof
wherein

R^2 is H, C_1 - C_{12} alkyl, C_3 - C_7 cycloalkyl, C_2 - C_8 alkenyl or C_2 - C_8 alkynyl, each optionally substituted with at least one radical selected from the group consisting of halogen, hydroxy, C_1 - C_4 alkoxy and C_1 - C_4 alkylthio; and

R^3 is H, C_1 - C_8 alkyl, C_1 - C_8 haloalkyl, C_2 - C_8 alkoxyalkyl or C_1 - C_6 hydroxyalkyl;

and



or an agriculturally useful salt thereof
wherein

- 5 R^4 is H, C_1-C_{12} alkyl, C_3-C_7 cycloalkyl, C_2-C_8 alkenyl or C_2-C_8 alkynyl, each optionally substituted with at least one radical selected from the group consisting of halogen, hydroxy, C_1-C_4 alkoxy and C_1-C_4 alkylthio.

This invention also relates to herbicidal compositions comprising effective amounts of the aforesaid mixtures and at least one of the following: surfactant, solid or liquid diluent. This invention also relates to a method of controlling undesired vegetation comprising
10 applying to the locus of the undesired vegetation herbicidally effective amounts of the aforesaid mixtures.

In the above recitations, the term "alkyl", used either alone or in compound words such as "haloalkyl" includes straight-chain or branched alkyl, such as, methyl, ethyl, *n*-propyl, *i*-propyl and the different butyl isomers. "Alkoxy" includes, for example, methoxy, ethoxy, *n*-propyloxy, isopropyloxy and the different butoxy isomers. The term "halogen", either
15 alone or in compound words such as "haloalkyl", includes fluorine, chlorine, bromine or iodine. Further, when used in compound words such as "haloalkyl", said alkyl may be partially or fully substituted with halogen atoms which may be the same or different. The total number of carbon atoms in a substituent group is indicated by the " C_i-C_j " prefix where
20 *i* and *j* are numbers from 1 to 12.

Some of the compounds of Formulae **IIa**, **IIb** and **IIc** can exist as one or more stereoisomers. The various stereoisomers include enantiomers, diastereomers, atropisomers and geometric isomers. One skilled in the art will appreciate that one stereoisomer may be more active and/or may exhibit beneficial effects when enriched relative to the other
25 stereoisomer(s) or when separated from the other stereoisomer(s). Additionally, the skilled artisan knows how to separate, enrich, and/or to selectively prepare said stereoisomers. Accordingly, the compounds of the invention may be present as a mixture of stereoisomers, individual stereoisomers, or as an optically active form.

Some of the compounds of Formulae **Ia** and **Ib** and Formulae **IIa**, **IIb** and **IIc** can be
30 present in the form of salts. In view of the close relationship between these compounds in their free forms and their agriculturally useful salts, including their equilibration under

physiological and environmental conditions, hereinbefore and hereinafter any reference to Formulae **Ia** and **Ib** and Formulae **IIa**, **IIb** and **IIc** is to be understood as including both the free compounds and their agriculturally useful salts, where appropriate and expedient.

The salts of the compounds of Formulae **Ia** and **Ib** and Formulae **IIa**, **IIb** and **IIc** include acid-addition salts with inorganic or organic acids such as hydrobromic, hydrochloric, nitric, phosphoric, sulfuric, acetic, butyric, fumaric, lactic, maleic, malonic, oxalic, propionic, salicylic, tartaric, 4-toluenesulfonic or valeric acids. The salts of the compounds of Formulae **Ia** and **Ib** and Formulae **IIa**, **IIb** and **IIc** also include those formed with organic bases (e.g., pyridine, ammonia, or triethylamine) or inorganic bases (e.g., hydrides, hydroxides, or carbonates of sodium, potassium, lithium, calcium, magnesium or barium) when the compound contains an acidic group such as a carboxylic acid or enol. Preferred salts include the lithium, sodium, potassium, triethylammonium, and quaternary ammonium salts.

Preferred for reasons of better weed control or improved crop safety are mixtures of the invention comprising:

- (1) the compound of Formula **Ia** and a compound selected from the compounds of Formula **IIa**, Formula **IIb** and Formula **IIc**; and
- (2) the compound of Formula **Ib** and a compound selected from the compounds of Formula **IIa**, Formula **IIb** and Formula **IIc**.

Most preferred for reasons of better weed control or improved crop safety are:

- (1a) the mixture of Preferred (1) comprising the compound of Formula **Ia** and the compound of Formula **IIa** where R^1 is CH_2CH_3 , which is ethyl 1-(2,4-dichlorophenyl)-5-(trichloromethyl)-1*H*-1,2,4-triazole-3-carboxylate (fenchlorazole-ethyl);
- (1b) the mixture of Preferred (1) comprising the compound of Formula **Ia** and the compound of Formula **IIb** where R^2 and R^3 are each CH_2CH_3 , which is diethyl 1-(2,4-dichlorophenyl)-4,5-dihydro-5-methyl-1*H*-pyrazole-3,5-dicarboxylate (mefenpyr-diethyl);
- (1c) the mixture of Preferred (1) comprising the compound of Formula **Ia** and the compound of Formula **IIc** where R^4 is $CH(CH_3)(CH_2)_4CH_3$, which is 1-methylhexyl [(5-chloro-8-quinolinyl)oxy]acetate (cloquintocet-mexyl);
- (2a) the mixture of Preferred (2) comprising the compound of Formula **Ib** and the compound of Formula **IIa** where R^1 is CH_2CH_3 , which is ethyl 1-(2,4-dichlorophenyl)-5-(trichloromethyl)-1*H*-1,2,4-triazole-3-carboxylate (fenchlorazole-ethyl);
- (2b) the mixture of Preferred (2) comprising the compound of Formula **Ib** and the compound of Formula **IIb** where R^2 and R^3 are each CH_2CH_3 , which is diethyl

1-(2,4-dichlorophenyl)-4,5-dihydro-5-methyl-1*H*-pyrazole-3,5-dicarboxylate (mefenpyr-diethyl); and


- (2c) the mixture of Preferred (2) comprising the compound of Formula **Ib** and the compound of Formula **Iic** where R^4 is $\text{CH}(\text{CH}_3)(\text{CH}_2)_4\text{CH}_3$, which is 1-methylhexyl [(5-chloro-8-quinolinyl)oxy]acetate (cloquintocet-mexyl).

Also preferred are herbicidal compositions comprising aforesaid preferred mixtures and at least one of the following: surfactant, solid or liquid diluent. Further preferred embodiments are methods of controlling undesired vegetation comprising applying to the locus of the vegetation herbicidally effective amounts of the aforesaid preferred mixtures and preferred herbicidal compositions.


The preferred crops for application of the mixtures of the invention are cool-season cereals including wheat, barley, oats and rye. Wheat and barley are particularly preferred crops.

DETAILS OF THE INVENTION

- The Formula **Ia** compound can be prepared as described in PCT publication WO 97/15576. The synthesis involves the reaction of the hydroxy compound of Formula 1 with diethylaminosulfur trifluoride (DAST) in dichloromethane solution.



- The Formula **Ib** compound can be prepared as described in U.S. 5,125,958. The synthesis involves coupling of the aniline of Formula 2 with ethyl acrylate (3) under diazotization conditions involving *tert*-butyl nitrite and copper(II) chloride in acetonitrile.



- Safeners of Formula **IIa** and their agriculturally useful salts can be prepared as described in U.S. 4,639,266. Safeners of Formula **IIb** and their agriculturally useful salts can be prepared by methods described in WO 91/07874. Safeners of Formula **IIc** and their agriculturally useful salts can be prepared by methods described in U.S. 4,881,966.

Combinations of the Formula **Ia** herbicide (Compound 1) and the Formula **Ib** herbicide (Compound 2) with the particular safeners of Formulae **IIa**, **IIb** and **IIc** identified in Index Table A are illustrative of the scope of this invention.

Index Table A

Safeners

<u>Safener</u>	<u>Formula</u>	<u>R¹</u>	<u>R²</u>	<u>R³</u>	<u>R⁴</u>
1	IIa	H	—	—	—
2	IIa	CH ₂ CH ₃	—	—	—
3	IIb	—	H	H	—
4	IIb	—	CH ₂ CH ₃	H	—
5	IIb	—	H	CH ₂ CH ₃	—
6	IIb	—	CH ₂ CH ₃	CH ₂ CH ₃	—
7	IIc	—	—	—	H
8	IIc	—	—	—	CH(CH ₃)(CH ₂) ₄ CH ₃

These combinations thus include those listed in the following Mixture Table A.

Mixture Table A

<u>Compound</u>	<u>Safener</u>	<u>Compound</u>	<u>Safener</u>	<u>Compound</u>	<u>Safener</u>
1	1	1	7	2	5
1	2	1	8	2	6
1	3	2	1	2	7
1	4	2	2	2	8
1	5	2	3		
1	6	2	4		

In Mixture Table A, the ratio of the herbicidal compound of Formula **Ia** (Compound 1) to each safener of Formulae **IIa**, **IIb** or **IIc** (Safeners 1–8) are in general 10:1 to 1:10 by weight, with weight ratios of 4:1 to 1:4 preferred for most uses. The ratio of the herbicidal compound of Formula **Ib** (Compound 2) to each safener of Formulae **IIa**, **IIb** or **IIc** (Safeners 1–8) are in general 10:1 to 1:10 by weight, with weight ratios of 4:1 to 1:4 preferred for most uses.

15 Formulation

The mixtures of at least one compound of Formulae **Ia** and **Ib** with the Formulae **IIa**, **IIb** or **IIc** compounds can be formulated in a number of ways:

- (a) the Formulae Ia and/or Ib and Formulae IIa, IIb or IIc compounds can be formulated separately and applied separately, with preferably the Formulae IIa, IIb or IIc safeners applied first; or
- (b) more preferably the Formulae Ia and/or Ib and Formulae IIa, IIb or IIc compounds can be applied simultaneously in an appropriate weight ratio, e.g., as a tank mix; or
- (c) most preferably the Formulae Ia and/or Ib and Formulae IIa, IIb or IIc compounds can be formulated together in the proper weight ratio.

Mixtures of at least one compound of Formulae Ia and Ib with the Formulae IIa, IIb or IIc compounds will generally be used in formulation with an agriculturally suitable carrier comprising a liquid or solid diluent and/or a surfactant wherein the formulation is consistent with the physical properties of the active ingredients, mode of application and environmental factors such as soil type, moisture and temperature. Useful formulations include liquids such as solutions (including emulsifiable concentrates), suspensions, emulsions (including micro-emulsions and/or suspoemulsions) and the like which optionally can be thickened into gels. Useful formulations further include solids such as dusts, powders, granules, pellets, tablets, films, and the like which can be water-dispersible ("wetable") or water-soluble. Active ingredients can be (micro)encapsulated and further formed into a suspension or solid formulation; alternatively the entire formulation of active ingredient can be encapsulated (or "overcoated"). Encapsulation can control or delay release of the active ingredients. Sprayable formulations can be extended in suitable media and used at spray volumes from about one to several hundred liters per hectare. High-strength compositions are primarily used as intermediates for further formulation.

The formulations will typically contain effective amounts of active ingredients, diluent and surfactant within the following approximate ranges which add up to 100 percent by weight.

	Weight Percent		
	<u>Active Ingredient</u>	<u>Diluent</u>	<u>Surfactant</u>
Water-Dispersible and Water-soluble Granules, Tablets and Powders.	5-90	0-94	1-15
Suspensions, Emulsions, Solutions (including Emulsifiable Concentrates)	5-50	40-95	0-15
Dusts	1-25	70-99	0-5
Granules and Pellets	0.01-99	5-99.99	0-15
High Strength Compositions	90-99	0-10	0-2

Typical solid diluents are described in Watkins, et al., *Handbook of Insecticide Dust Diluents and Carriers*, 2nd Ed., Dorland Books, Caldwell, New Jersey. Typical liquid diluents are described in Marsden, *Solvents Guide*, 2nd Ed., Interscience, New York, 1950. *McCutcheon's Detergents and Emulsifiers Annual*, Allured Publ. Corp., Ridgewood, New Jersey, as well as Sisely and Wood, *Encyclopedia of Surface Active Agents*, Chemical Publ. Co., Inc., New York, 1964, list surfactants and recommended uses. All formulations can contain minor amounts of additives to reduce foam, caking, corrosion, microbiological growth and the like, or thickeners to increase viscosity.

Surfactants include, for example, polyethoxylated alcohols, polyethoxylated alkylphenols, polyethoxylated sorbitan fatty acid esters, dialkyl sulfosuccinates, alkyl sulfates, alkylbenzene sulfonates, organosilicones, *N,N*-dialkyltaurates, lignin sulfonates, naphthalene sulfonate formaldehyde condensates, polycarboxylates, and polyoxyethylene/polyoxypropylene block copolymers. Solid diluents include, for example, clays such as bentonite, montmorillinite, attapulgite and kaolin, starch, sugar, silica, talc, diatomaceous earth, urea, calcium carbonate, sodium carbonate and bicarbonate, and sodium sulfate. Liquid diluents include, for example, water, *N,N*-dimethylformamide, dimethyl sulfoxide, *N*-alkylpyrrolidone, ethylene glycol, polypropylene glycol, paraffins, alkylbenzenes, alkyl naphthalenes, oils of olive, castor, linseed, tung, sesame, corn, peanut, cotton-seed, soybean, rape-seed and coconut, fatty acid esters, ketones such as cyclohexanone, 2-heptanone, isophorone and 4-hydroxy-4-methyl-2-pentanone, and alcohols such as methanol, cyclohexanol, decanol and tetrahydrofurfuryl alcohol.

Solutions, including emulsifiable concentrates, can be prepared by simply mixing the ingredients. Dusts and powders can be prepared by blending and, usually, grinding as in a hammer mill or fluid-energy mill. Suspensions are usually prepared by wet-milling; see, for example, U.S. 3,060,084. Granules and pellets can be prepared by spraying the active material upon preformed granular carriers or by agglomeration techniques. See Browning, "Agglomeration", *Chemical Engineering*, December 4, 1967, pp 147-48, *Perry's Chemical Engineer's Handbook*, 4th Ed., McGraw-Hill, New York, 1963, pages 8-57 and following, and WO 91/13546. Pellets can be prepared as described in U.S. 4,172,714. Water-dispersible and water-soluble granules can be prepared as taught in U.S. 4,144,050, U.S. 3,920,442 and DE 3,246,493. Tablets can be prepared as taught in U.S. 5,180,587, U.S. 5,232,701 and U.S. 5,208,030. Films can be prepared as taught in GB 2,095,558 and U.S. 3,299,566.

For further information regarding formulation, see U.S. 3,235,361, Col. 6, line 16 through Col. 7, line 19 and Examples 10-41; U.S. 3,309,192, Col. 5, line 43 through Col. 7, line 62 and Examples 8, 12, 15, 39, 41, 52, 53, 58, 132, 138-140, 162-164, 166, 167 and 169-182; U.S. 2,891,855, Col. 3, line 66 through Col. 5, line 17 and Examples 1-4;

Klingman, *Weed Control as a Science*, John Wiley and Sons, Inc., New York, 1961, pp 81–96; and Hance et al., *Weed Control Handbook*, 8th Ed., Blackwell Scientific Publications, Oxford, 1989.

In the following Examples, all percentages are by weight and all formulations are prepared in conventional ways. Compound 1 is the compound of Formula **Ia**, which is (6*S*-*cis*)-1-chloro-*N*-[2-chloro-4-fluoro-5-(6-fluorotetrahydro-1,3-dioxo-1*H*-pyrrolo-[1,2-*c*]imidazol-2(3*H*)-yl)phenyl]methanesulfonamide, and Compound 2 is the compound of Formula **Ib**, which is ethyl α ,2-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1*H*-1,2,4-triazol-1-yl]-4-fluorobenzenepropanoate (carfentrazone-ethyl). Safener numbers are defined in Index Table A (page 6).

Example A

High Strength Concentrate

	Compound 1	9.0%
	Safener 2	89.5%
15	silica aerogel	0.5%
	synthetic amorphous fine silica	1.0%.

Example B

Wettable Powder

	Compound 1	52.0%
20	Safener 6	13.0%
	dodecylphenol polyethylene glycol ether	2.0%
	sodium ligninsulfonate	4.0%
	sodium silicoaluminate	6.0%
	montmorillonite (calcined)	23.0%.

Example C

Granule

	Compound 1	5.0%
	Safener 8	5.0%
	attapulgate granules (low volatile matter, 0.71/0.30 mm; U.S.S. No. 25–50 sieves)	90.0%.
30		

Example DAqueous Suspension

	Compound 1	5.0%
	Safener 8	20.0%
5	hydrated attapulgite	3.0%
	crude calcium ligninsulfonate	10.0%
	sodium dihydrogen phosphate	0.5%
	water	61.5%.

Example E10 Extruded Pellet

	Compound 1	2.3%
	Safener 8	22.7%
	anhydrous sodium sulfate	10.0%
	crude calcium ligninsulfonate	5.0%
15	sodium alkyl naphthalenesulfonate	1.0%
	calcium/magnesium bentonite	59.0%.

Example FHigh Strength Concentrate

	Compound 2	9.0%
20	Safener 2	89.5%
	silica aerogel	0.5%
	synthetic amorphous fine silica	1.0%.

Example GWettable Powder

25	Compound 2	52.0%
	Safener 6	13.0%
	dodecylphenol polyethylene glycol ether	2.0%
	sodium ligninsulfonate	4.0%
	sodium silicoaluminate	6.0%
30	montmorillonite (calcined)	23.0%.

Example HHigh Strength Concentrate

	Compound 2	49.2%
	Safener 8	49.3%
5	silica aerogel	0.5%
	synthetic amorphous fine silica	1.0%.

Utility

Test results indicate that combinations of the chlorofluorobenzene herbicides of Formulae **Ia** and/or **Ib** with a safener of Formulae **IIa**, **IIb** or **IIc** affords highly active herbicidal mixtures, providing unexpected safening on cool-season cereal crops such as wheat, barley and rye, while retaining good efficacy for controlling many problem weeds in these crops. Certain combinations even show a greater-than-additive or synergistic effect on selected weeds.

Herbicidally effective amounts of the compounds of Formulae **Ia** and/or **Ib** and antidotally effective amounts of the safeners of Formulae **IIa**, **IIb** or **IIc** will vary depending on environmental conditions, formulation, method of application, amount and type of vegetation present, etc. The use rate ratios of Formula **Ia** to Formulae **IIa**, **IIb** or **IIc** are in general 10:1 to 1:10 by weight, with weight ratios of 4:1 to 1:4 preferred for most uses. The use rate ratios of Formula **Ib** to Formulae **IIa**, **IIb** or **IIc** are in general 10:1 to 1:10 by weight, with weight ratios of 4:1 to 1:4 preferred for most uses. In general, the Formula **Ia** compound is applied at a rate from 1 to 100 g ai/ha, and preferably the Formula **Ia** compound is applied at a rate from 2 to 20 g ai/ha. In general, the Formula **Ib** compound is applied at a rate from 1 to 100 g ai/ha, and preferably the Formula **Ib** compound is applied at a rate from 8 to 40 g ai/ha. In general, the Formulae **IIa**, **IIb** or **IIc** compounds are applied at rates from 0.1 to 1000 g ai/ha, and preferably the Formulae **IIa**, **IIb** or **IIc** compounds are applied at rates from 0.5 to 120 g ai/ha.

For selective control of certain weed species in cereal crops, the mixtures of the invention are best applied when the cereal plants are at about the 1-leaf through tillering stage and the weed plants are correspondingly young. One skilled in the art can readily determine application rates and ratios of the herbicides of Formulae **Ia** and/or **Ib** to the safeners of Formulae **IIa**, **IIb** or **IIc** as well as timing necessary for the desired level of weed control and crop safety.

The Formulae **Ia** and/or **Ib** and Formulae **IIa**, **IIb** or **IIc** mixtures of this invention can additionally be used in combination with other commercial herbicides, insecticides or fungicides. A mixture of one or more of the following herbicides with the Formulae **Ia** and/or **Ib** and Formulae **IIa**, **IIb** or **IIc** mixtures of this invention may be particularly useful

for weed control. Examples of other herbicides particularly useful as mixture partners are: 2,4-D, bentazon, benzoylpropethyl, bromoxynil, chlorsulfuron, chlortoluron, clodinafop, clopyralid, cyanazine, diallate, dicamba, diclofop, difenzoquat, diflufenican, fenchlorazole, fenoxaprop, flamprop-M-isopropyl, fluroglycophen-ethyl, fluroxypyr, glyphosate, imazamethabenz, ioxynil, isoproturon, linuron, MCPA, mecoprop, methabenzthiazuron, metosulam, metoxuron, metribuzin, metsulfuron methyl, *N*-[[[(4,6-methoxy-2-pyrimidinyl)-amino]carbonyl]-2-(ethylsulfonyl)imidazo[1,2-*a*]pyridine-3-sulfonamide (ISO-proposed: sulfosulfuron), neburon, paraquat, pendimethalin, propanil, prosulfocarb, prosulfuron, quinclorac, simazine, terbutryn, thifensulfuron methyl, tralkoxydim, triallate, triasulfuron, tribenuron methyl and trifluralin.

In certain instances, combinations with other herbicides having a similar spectrum of control but a different mode of action will be particularly advantageous for resistance management.

The following data demonstrate the surprising safening of the mixtures of this invention on cool-season cereal crops while retaining excellent efficacy against many weeds. The weed control afforded by the mixtures of this invention are not limited, however, to these species. Compound 1 is the compound of Formula **Ia**, which is (6*S*-*cis*)-1-chloro-*N*-[2-chloro-4-fluoro-5-(6-fluorotetrahydro-1,3-dioxo-1*H*-pyrrolo[1,2-*c*]imidazol-2(3*H*)-yl)phenyl]methanesulfonamide, and Compound 2 is the compound of Formula **Ib**, which is ethyl α ,2-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1*H*-1,2,4-triazol-1-yl]-4-fluorobenzenepropanoate (carfentrazone-ethyl). Safener numbers are defined in Index Table A (page 6).

TEST A

Test Protocol

The crops: winter wheat (*Triticum aestivum*, TRZAW), spring wheat (*Triticum aestivum*, TRZAS) and spring barley (*Hordeum vulgare*, HORVS); and the weeds: pigweed (*Amaranthus retroflexus*, AMARE), common lambsquarters (*Chenopodium album*, CHEAL), cleavers (*Galium aparine*, GALAP), kochia (*Kochia scoparia*, KCHSC), henbit (*Lamium amplexicaule*, LAMAM), scentless chamomile (*Matricaria inodora*, MATIN), wild buckwheat (*Polygonum convolvulus*, POLCO), Russian thistle (*Salsola kali*, SASKR), wild mustard (*Sinapsis arvensis*, SINAR), black nightshade (*Solanum nigrum*, SOLNI) and common chickweed (*Stellaria media*, STEME) were planted in a sterilized growing medium at the depths indicated and grown in a greenhouse approximately 10–17 days to the stages shown:

Species	Depth (cm)	Growth Stage	Species	Depth (cm)	Growth Stage
winter wheat	2.5	1-2 leaf	<i>L. amplexicaule</i>	0.4	1-2 leaf pair
spring wheat	2.5	1-2 leaf	<i>M. inodora</i>	0.4	2-4 leaf
spring barley	2.5	1-2 leaf	<i>P. convolvulus</i>	1.2	1-2 leaf
<i>A. retroflexus</i>	0.4	1-2 leaf	<i>S. kali</i>	1.8	2-4 leaf
<i>C. album</i>	0.5	2-4 leaf	<i>S. arvensis</i>	1.2	2-3 leaf
<i>G. aparine</i>	1.2	2-3 whorl	<i>S. nigrum</i>	1.8	1 leaf
<i>K. scoparia</i>	0.4	4-8 leaf	<i>S. media</i>	0.4	2-3 leaf pair

The pots were irrigated as needed with water containing 200 ppm N from 20-20-20 fertilizer. The greenhouse was maintained at approximately 21-24 °C during the day and 13-16 °C at night with 14 hours of daylight supplemented as needed by metal halide lamps.

Compound 1 was applied at 8 g ai/ha postemergence alone and in combination with 1, 2 and 4 g ai/ha of Safener 2, Safener 6 or Safener 8. All samples were sprayed in a non-phytotoxic solvent in a volume of 309 L/ha. Assessments of crop response and weed control were made by visual inspection 14 days following application. A visual rating system was used based on a percentage scale from 0 to 100%, relative to adjacent untreated control plants. On this scale 0% represents no visual differences relative to an untreated control, and 100% represents complete kill of the given crop or weed species.

Table A represents visual assessments of crop response and specific weed control with Compound 1 applied alone and in combination with Safener 2, Safener 6 or Safener 8. Different ratios of Compound 1 to Safener 2, ratios of Compound 1 to Safener 6, ratios of Compound 1 to Safener 8, and different formulation types, also provide useful crop safening and weed control from the combinations.

TABLE A

Effect of Compound 1 and as Active Ingredient Alone
and in Mixtures with Safeners 2, 6 and 8

Application Rates (g ai/ha)		Crops			Weeds				
		T	T	H	A	C	G	K	L
		R	R	O	M	H	A	C	A
		Z	Z	R	A	E	L	H	M
		A	A	V	R	A	A	S	A
Compound 1	Safener	W	S	S	E	L	P	C	M
<i>Compound 1 Alone</i>									
8	—	10	10	10	100	100	80	100	65
<i>With Safener 2</i>									
8	1	5	5	5	100	100	90	100	65
8	2	5	5	10	100	100	100	100	80
8	4	5	5	2	100	100	100	100	60
<i>With Safener 6</i>									
8	1	5	5	5	100	100	85	100	85
8	2	2	5	5	100	100	80	100	85
8	4	2	5	5	100	100	98	100	65
<i>With Safener 8</i>									
8	1	2	5	5	100	100	100	100	70
8	2	2	5	5	100	100	100	100	65
8	4	2	5	2	100	100	100	100	65

TABLE A continued

Application Rates (g ai/ha)		Weeds					
		M A T I N	P O L C O	S A S K R	S I N A R	S O L N I	S T E M E
Compound 1	Safener						
<i>Compound 1 Alone</i>							
8	—	100	100	100	100	100	80
<i>With Safener 2</i>							
8	1	100	100	100	100	100	100
8	2	100	100	100	85	100	100
8	4	100	100	100	100	100	100
<i>With Safener 6</i>							
8	1	100	100	100	100	100	100
8	2	100	100	100	100	100	100
8	4	100	100	100	100	100	100
<i>With Safener 8</i>							
8	1	100	100	100	100	100	100
8	2	100	100	100	100	100	100
8	4	100	100	100	100	100	100

As can be seen from Table A, mixtures of Compound 1 with Safener 2, Safener 6 or Safener 8 were found to cause significantly less injury to the wheat and barley compared to Compound 1 applied alone. Compound 1 alone and in combination with Safener 2, Safener 6 or Safener 8 gave 100% control of most weed species. In this test, combinations of Compound 1 with Safener 2 or Safener 8 appeared to increase control of *Galium aparine*. The surprising combination of crop safening and excellent weed control make these mixtures particularly commercially valuable.

TEST B

10 Test Protocol

The crops: winter wheat (*Triticum aestivum*, TRZAW), spring wheat (*Triticum aestivum*, TRZAS), winter barley (*Hordeum vulgare*, HORVW) and spring barley (*Hordeum vulgare*, HORVS); and the weeds: common lambsquarters (*Chenopodium album*, CHEAL), cleavers (*Galium aparine*, GALAP), henbit (*Lamium amplexicaule*, LAMAM), wild

buckwheat (*Polygonum convolvulus*, POLCO), Russian thistle (*Salsola kali*, SASKR), and black nightshade (*Solanum nigrum*, SOLNI) were planted in a sterilized growing medium at the depths indicated and grown in a greenhouse approximately 10–17 days to the stages shown:

Species	Depth (cm)	Growth Stage	Species	Depth (cm)	Growth Stage
winter wheat	2.5	1–2 leaf	<i>G. aparine</i>	1.2	2–3 whorl
spring wheat	2.5	1–2 leaf	<i>L. amplexicaule</i>	0.4	1–2 leaf pair
winter barley	2.5	1–2 leaf	<i>P. convolvulus</i>	1.2	1–2 leaf
spring barley	2.5	1–2 leaf	<i>S. kali</i>	1.8	2–4 leaf
<i>C. album</i>	0.5	2–4 leaf	<i>S. nigrum</i>	1.8	1 leaf

- 5 The pots were irrigated as needed with water containing 200 ppm N from 20-20-20 fertilizer. The greenhouse was maintained at approximately 21–24 °C during the day and 13–16 °C at night with 14 hours of daylight supplemented as needed by metal halide lamps.

Test B was comprised of six replicates. Compound 2 was applied at 16 g ai/ha postemergence alone and in combination 8 g ai/ha of Safener 2, Safener 6 or Safener 8. All
 10 samples were sprayed in a non-phytotoxic solvent in a volume of 309 L/ha. Assessments of crop response and weed control were made by visual inspection 5 days following application. A visual rating system was used based on a percentage scale from 0 to 100%, relative to adjacent untreated control plants. On this scale 0% represents no visual differences relative to an untreated control, and 100% represents complete kill of the given crop or weed species.

15 Table B represents mean visual assessments of crop response and specific weed control with Compound 2 applied alone and in combination with Safener 2, Safener 6 or Safener 8. Different ratios of Compound 2 to Safener 2, ratios of Compound 2 to Safener 6, ratios of Compound 2 to Safener 8, and different formulation types, also provide useful crop safening and weed control from the combinations.

TABLE B

Mean* Effect of Compound 2 and as Active Ingredient Alone
and in Mixtures with Safeners 2, 6 and 8

Application Rates (g ai/ha)		Crops				Weeds				
		H	T	T	H	G	L	S	S	P
		O	R	R	O	A	A	A	O	O
		R	Z	Z	R	L	M	S	L	L
		V	A	A	V	A	A	K	N	C
Compound 2	Safener	W	W	S	S	P	M	R	I	O
Compound 2 alone										
16	—	9	9	22	8	100	100	100	100	100
With Safener 2										
16	8	8	6	13	3	99	100	98	100	100
With Safener 6										
16	8	6	4	5	2	100	100	99	100	100
With Safener 8										
16	8	4	4	8	3	100	100	100	100	100

* Values are the mean response of six replicates.


- 5 As can be seen from Table B, mixtures of Compound 2 with Safener 2, Safener 6 or Safener 8 were found to cause significantly less injury to wheat and barley compared to Compound 2 applied alone. Compound 2 alone and in combination with Safener 2, Safener 6 or Safener 8 gave 98–100% control of the weed species. The surprising combination of crop safening and excellent weed control make these mixtures particularly commercially
- 10 valuable.

18


CLAIMS

What is claimed is:

1. A mixture comprising a herbicidally effective amount of at least one herbicidal compound selected from the group consisting of:


**Ia**

which is (6*S*-*cis*)-1-chloro-*N*-[2-chloro-4-fluoro-5-(6-fluorotetrahydro-1,3-dioxo-1*H*-pyrrolo-[1,2-*c*]imidazol-2(3*H*)-yl)phenyl]methanesulfonamide,
or an agriculturally useful salt thereof; and

**Ib**


which is ethyl α,2-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1*H*-1,2,4-triazol-1-yl]-4-fluorobenzenepropanoate,
or an agriculturally useful salt thereof;

in combination with an antidotally effective amount of a safener compound selected from the group consisting of:

**IIa**

or an agriculturally useful salt thereof
wherein


R^1 is C_1 - C_{12} alkyl, C_3 - C_7 cycloalkyl, C_2 - C_8 alkenyl or C_2 - C_8 alkynyl, each optionally substituted with at least one radical selected from the group consisting of halogen, hydroxy, C_1 - C_4 alkoxy and C_1 - C_4 alkylthio;



or an agriculturally useful salt thereof

wherein

- 5 R^2 is H, C_1-C_{12} alkyl, C_3-C_7 cycloalkyl, C_2-C_8 alkenyl or C_2-C_8 alkynyl, each optionally substituted with at least one radical selected from the group consisting of halogen, hydroxy, C_1-C_4 alkoxy and C_1-C_4 alkylthio; and
- R^3 is H, C_1-C_8 alkyl, C_1-C_8 haloalkyl, C_2-C_8 alkoxyalkyl or C_1-C_6 hydroxyalkyl; and



- 10 or an agriculturally useful salt thereof

wherein

- R^4 is H, C_1-C_{12} alkyl, C_3-C_7 cycloalkyl, C_2-C_8 alkenyl or C_2-C_8 alkynyl, each optionally substituted with at least one radical selected from the group consisting of halogen, hydroxy, C_1-C_4 alkoxy and C_1-C_4 alkylthio.
- 15 2. A mixture of Claim 1 wherein the herbicidal compound is (6*S*-*cis*)-1-chloro-*N*-[2-chloro-4-fluoro-5-(6-fluorotetrahydro-1,3-dioxo-1*H*-pyrrolo[1,2-*c*]imidazol-2(3*H*)-yl)phenyl]methanesulfonamide.
3. The mixture of Claim 2 wherein the safener compound is ethyl 1-(2,4-dichlorophenyl)-5-(trichloromethyl)-1*H*-1,2,4-triazole-3-carboxylate.
- 20 4. The mixture of Claim 2 wherein the safener compound is diethyl 1-(2,4-dichlorophenyl)-4,5-dihydro-5-methyl-1*H*-pyrazole-3,5-dicarboxylate.
5. The mixture of Claim 2 wherein the safener compound is 1-methylhexyl [(5-chloro-8-quinolinyl)oxy]acetate.

6. A mixture of Claim 1 wherein the herbicidal compound is ethyl α ,2-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1*H*-1,2,4-triazol-1-yl]-4-fluorobenzene-propanoate.

7. The mixture of Claim 6 wherein the safener compound is ethyl
5 1-(2,4-dichlorophenyl)-5-(trichloromethyl)-1*H*-1,2,4-triazole-3-carboxylate.

8. The mixture of Claim 6 wherein the safener compound is diethyl
1-(2,4-dichlorophenyl)-4,5-dihydro-5-methyl-1*H*-pyrazole-3,5-dicarboxylate.

9. The mixture of Claim 6 wherein the safener compound is 1-methylhexyl
[(5-chloro-8-quinolinyl)oxy]acetate.

10 10. An agriculturally suitable composition for controlling the growth of undesired
vegetation comprising an effective amount of the mixture of any of Claims 1–9 and at least
one of the following: surfactant, solid or liquid diluent.

11. A method for controlling the growth of undesired vegetation by applying to the
locus of the undesired vegetation a herbicidally effective amount of the mixture of any of
15 Claims 1–9.

12. The method of Claim 11 wherein the locus of the undesired vegetation is a crop
of wheat or barley.

INTERNATIONAL SEARCH REPORT

International Application No

PC./US 99/10359

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A01N43/90 A01N43/653 //(A01N43/90,43:653,43:56,43:42),
(A01N43/653,43:653,43:56,43:42)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 11574 A (CIBA GEIGY AG ;GLOCK JUTTA (CH); HUDETZ MANFRED (US); KERBER ELMAR) 25 April 1996 (1996-04-25) see page 1, par. 3 and 4, and safeners (IIa), (IIb) and (IIb1) (Tables 16 - 18). ----	1-12
A	EP 0 492 367 A (HOECHST AG) 1 July 1992 (1992-07-01) page 6, line 15 - page 8, line 57 ----	1-12
A	WO 91 07874 A (HOECHST AG) 13 June 1991 (1991-06-13) cited in the application page 6, line 32 - page 7, line 6 -----	1-12

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

° Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance
 "E" earlier document but published on or after the international filing date
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 "O" document referring to an oral disclosure, use, exhibition or other means
 "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
 "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
 "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
 "&" document member of the same patent family

Date of the actual completion of the international search

8 September 1999

Date of mailing of the international search report

15/09/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Klaver, J

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/10359

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9611574 A	25-04-1996	AU 3653695 A	06-05-1996
		BR 9509374 A	30-12-1997
		CA 2199636 A	25-04-1996
		EP 0786937 A	06-08-1997
		JP 10507189 T	14-07-1998
		PL 319092 A	21-07-1997
		ZA 9508712 A	10-07-1996
EP 0492367 A	01-07-1992	AU 8989791 A	25-06-1992
		CA 2058284 A	22-06-1992
		HU 211899 B	29-01-1996
		JP 4308508 A	30-10-1992
		MX 9102750 A	01-06-1992
		US 5488027 A	30-01-1996
WO 9107874 A	13-06-1991	DE 3939503 A	06-06-1991
		AT 163124 T	15-02-1998
		AU 653506 B	06-10-1994
		AU 6886391 A	26-06-1991
		CA 2069901 A	31-05-1991
		CN 1052115 A	12-06-1991
		DE 59010806 D	19-03-1998
		EP 0635996 A	01-02-1995
		ES 2114862 T	16-06-1998
		GR 3026248 T	29-05-1998
		IL 96496 A	29-12-1994
		JP 5503086 T	27-05-1993
		LT 711 A, B	31-01-1995
		LV 10359 A, B	20-02-1995
		US 5700758 A	23-12-1997
		US 5703008 A	30-12-1997